

Veneer Zement
Ciment pour facette
Cemento para Veneer
Cemento per faccette
Fasadcement
Veneer/facade cement
Laminaattisementti
Veneercement
Sementering av laminater
Cimento para Facetas
КпнЯб гйб ьшейт
Цемент Для Облицовок



CONTENTS:

2 Syringes CHOICE 2 Shade: TRANSLUCENT (4.5g ea.), 1 Syringe CHOICE 2 Shade: A1 (4.5g), 1 Syringe CHOICE 2 Shade: B1 (4.5g), 1 Syringe CHOICE 2 Shade: MILKY BRIGHT (4.5g), 1 Syringe UNI-ETCH® w/BAC (5g), 1 Syringe 9.5% Porcelain Etchant (5g), 1 Bottle each Bis-Silane™ Parts A & B (6ml ea.), 1 Bottle each ALL-BOND 3® Parts A & B (6ml ea.), 1 Bottle Porcelain Bonding Resin (6ml), Accessories, Instructions



BISCO, Inc.
1100 W. Irving Park Rd.
Schaumburg, IL 60193
U.S.A.
1-847-534-6000
www.bisco.com



















https://www.bisco.com/symbols-glossary/



Choice™ 2 PORCELAIN VENEER RESTORATION PREPARATION & CEMENTATION

PORCELAIN VENEER RESTORATION PREPARATION

If the veneer has not been etched by the laboratory, etch with 9.5% PORCELAIN ETCHANT for 90 seconds.



Rinse with water.



Thoroughly air dry.





Mix BIS-SILANE[™] Parts A & B and apply 1-2 coats to the internal surface. Wait for 30 seconds.



Dry with air syringe.

TOOTH PREPARATION FOR PORCELAIN VENEER

Clean the preparation with a slurry of pumice. Rinse and dry. Etch dentin and enamel using UNI-ETCH® for 15 seconds.



Rinse thoroughly.



Lightly air dry. Leave preparation visibly moist.




Mix ALL-BOND 3® Parts A & B and apply 1-2 consecutive coats to the dentin and enamel.

Note: Additional coats may be required if a lot of dentin is exposed.



Air dry for a minimum of 10 seconds to evaporate solvent.



Light cure for 10 seconds

PORCELAIN VENEER CEMENTATION


Apply a very thin layer of PORCELAIN BONDING RESIN to the inside of the veneer. DO NOT LIGHT CURE.



Line inside of veneer with CHOICE™ 2 Veneer Cement



Working in pairs, gently seat one veneer at a time.



Light cure each for 3-5 seconds to tack the veneers into place. Remove excess cement. Light cure each veneer for 40 seconds from lingual and 40 seconds from the facial.





