

Instructions for Use



BISCO, Inc.
1100 W. Irving Park Road
Schaumburg, IL 60193
U.S.A.

1-847-534-6000

1-800-247-3368

IN-204R6

Rev. 9/18

Caution: U.S. Federal law restricts this device to sale by or on the order of a licensed Dentist.

*The **CEMENT** kit containing etchant w/BAC is not CE marked.

eCEMENT®

Adhesive Cementation System for Lithium Disilicate Restorations

GENERAL INFORMATION

CEMENT is a resin cementation system designed to simplify the placement of Lithium Disilicate ceramic (e.g. IPS e.max®) restorations. Lithium Disilicate is a silica-based glass ceramic available in pressed and fused forms, with both high and medium opacities, and high and low translucencies. The system contains a lignite, and can be used for all types of restorative procedures. All-BOND UNIVERSAL restorative will not etch prior to application. The **CEMENT** system may also be used for cementation of other silica-based glass ceramic restorations. **CEMENT TRY-IN PASTES** are available separately and may be used to visualize and pre-determine the aesthetics of the final cementation, they are water-soluble and rinse easily.

Indications for Use:

• **CEMENT** LC: thin restorations with high translucency, where a light cured only cement is preferred (includes veneers and crowns and inlays). The system contains a lignite, and can be used for all types of restorative procedures.

• **CEMENT** DC: Thicker restorations with medium or high opacity and low translucency, where a dual-cured cement is required (includes opaque veneers, inlays, onlays and crowns, and free-unit bridges, primarily in the posterior region).

NOTE: Includes high strength zirconia/alumina restorations.

Contraindications:

• Do not use ALL-BOND UNIVERSAL® on patients who have a history of severe allergic reactions to methacrylate resins.

Warnings:

• ALL-BOND UNIVERSAL and **PORECLAIN PRIMER** is highly flammable.

• Do not use this product until you have thoroughly read and understood these instructions. Hydrofluoric acids is an aggressive chemical if it is severe eye and tissue irritant. If accidentally splashed into the eye, flush with copious amounts of water for 15 minutes and seek immediate medical attention.

• Hydrofluoric acid also releases hydrogen fluoride gas which is extremely dangerous to health.

The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

• Do NOT use PORCELAIN ETCHANT® (4% Hf) to etch exposed dentin or enamel surfaces.

• Verify flow of etchants prior to applying intracanal. If needed, replace tip and re-check. Only use recommended tips.

• Re-etch after rinsing with the tooth cap.

• Avoid splashing into the eyes. ALL-BOND UNIVERSAL comes into contact with the eyes, flush with copious amounts of water and seek medical attention.

• The operator and assistant must wear protective gloves. Rubber dam isolation must be used for intracanal repair.

